

# Diagnostic informativity of the volume MDCT-angiography and MR-cholangiography in the pre- and intraoperative periods for the examination of donors of a liver fragment

Feruz Gafurovich NAZIROV, Nigora Muxsumovna DJURAEVA, Nargiza Tulkinovna VAKHIDOVA, Oybek Avaxzanovich OMONOV and Umid Ravshanovich SALIMOV ✉✉

Republican Specialized Scientific and Practical Medical Center of Surgery named after academician V.Vakhidov Tashkent, Uzbekistan

✉ Corresponding author's Email: sidikov@yandex.ru

## ABSTRACT

**Introduction.** The first transplantation of a liver fragment from a living donor was performed in Uzbekistan on February 12, 2018 at the Republican Specialized Scientific Practical Medical Center of Surgery named after acad.V. Vakhidov. This event laid the foundation for a new direction for domestic clinical practice that meets the current level of world medicine development. **Aim.** The aim of the study was to determine the diagnostic information content of preoperative data of the volume multi detector computed tomography (MDCT) angiography and magnetic resonance cholangiography (MRCG) when compared with intraoperative ones at examining related donors for liver fragment transplantation (LFT). **Methods.** Total of 88 potential donors of a liver fragment aged from 19 to 58 years (53 men and 35 women) were examined for the period 2017-2019. Sixteen donors were undergone liver resection to obtain a transplant: the right lobe of the livers in 12 people and the left lobe in 4 people. **Results.** Compared with intraoperative data, the main arteries supplying the transplant planned for resection were identified with MDCT-angiography in 98.4% of cases ( $P<0.05$ ). Variations of the portal bed according to MDCT-angiography in comparison with intraoperative ones were determined in 93.8% of cases ( $P<0.05$ ). Intraoperatively revealed the main trunks of the venous outflow were determined by MDCT-angiography in 95.7% of cases ( $P<0.05$ ). **Recommendation.** We suggest that MDCT angiography and MRCG is a highly informative and important method for estimation the condition of the liver in transplant planning.

## Original Article

PII: S225199391900024-9

Rec. 11 October 2019

Rev. 20 November 2019

Pub. 25 November 2019

## Keywords

Liver transplant,  
Contrast agent,  
MDCT-angiography,  
Magnetic resonance  
cholangiography.

## INTRODUCTION

The undoubted advantage of receiving a transplant from a living donor in the planned preparation of related donors and the possibility of a qualitative choice of donor hepatic parenchyma is an important factor for a favorable prognosis of the surgery's outcome. Negative hemodynamic and drug effects are practically excluded for the donor in the case of a related transplantation at the preoperative stage [1-4]. The indisputable advantage of a liver fragment transplantation (LFT) from a living related donor is independence from the system of providing cadaveric organs, which means that planning the timing of the operation depending on the state of the recipient [5-10].

In contrast to cadaveric liver transplantation, the use of an organ fragment from a close relative allows to count on its more favorable immunological adaptation in the recipient's body [11-13]. The first transplantation of a liver fragment from a living related donor was performed in Uzbekistan on February 12, 2018 at the Republican Specialized Scientific Practical Medical Center of Surgery (RSSPMCS) named after acad. V.Vakhidov. This event laid the foundation for a new direction for domestic clinical practice that meets the current level of world medicine development [12, 14].

All main disadvantages of related liver transplantation are associated with a potential risk to the health and life of the donor which causes certain skepticism and ethical questions among the general medical community. Thereby, one of the important aspects of planning the surgery of related donors is a detailed examination of the selected potential donor [11, 15-20]. After conducting clinical and laboratory studies instrumental examination methods are performed at the final stage, which includes volume MDCT angiography and magnetic resonance cholangiography (MRCG). The main task is to clarify the technical capabilities of obtaining a full-fledged liver transplant while fully preserving the possibilities of donor rehabilitation. It is necessary to determine the following for doing this: 1) the size and weight of the donor liver fragment to be removed and the remaining part (liver stump); 2) an anatomical version of the circulatory system structure

entering the porta of the donor liver, venous outflow from the organ and architectonics of the biliary system [13, 21-24]. Since its introduction to the clinics, MDCT angiography has become the main method of instrumental examination of related donors of a liver fragment [2, 6, 16, 25, 26]. In accordance with the requirements of transplant doctors, a special MDCT scan protocol (all in one) was developed, which includes four phases of the study: native, arterial, portal and venous [27-31]. To determine the diagnostic information content of the preoperative results of volume MDCT angiography and MRCG in the rebound of potential liver donors, as well as their comparison with the intraoperative anatomical picture.

## MATERIAL AND METHODS

### Ethical approval

The review board and ethics committee of RSCS named after acad. V.Vakhidov approved the study protocol and informed consents were taken from all the participants.

Total of 88 potential donors of a liver fragment – 53 men and 35 women – aged from 19 to 58 years (the mean age was 34.1±9.06 years) were examined. All patients were examined in RSSPMCS named after acad.V. Vakhidov for the period 2017-2019. At the time of the present study, 16 donors were undergone liver resection to obtain a transplant: in 12 people (the right lobe, RL), in 4 people (the left lobe of the liver, LL). All related donors underwent four-phase MDCT angiography and MRCG of liver. The investigations were carried out on the wide-detector MDCT (Aquilion One - 640 Genesis, Cannon, Japan), as well as MRI on the Signa HDxt with a magnetic field of 1.5 T (GE). The rate of contrast agent (CA) introduction was 5.0–5.5 ml/sec, the amount of CA varied 100±10 ml (Unigexol 350).

MDCT investigation began with an inserting of the catheter into the external cubital vein followed by the placement and instruction of the patient about the progress of the upcoming procedure. An automatic syringe-injector of the “Ulrich” company was used for injection, having previously collected 100-120 ml of an iodine-containing contrast preparation into the flask for injection with an iodine concentration of 350-370 mg / ml and an injection rate of 4.5-5.0 ml/s. The selection protocol in this case was a multiphase scanning protocol with selected parameters, which included a native study, arterial, portal and delayed phases (phase of the hepatic veins) scans performed on one breath hold in each phase (Table 1). Magnetic resonance imaging (MRI) investigation of the patient began with laying on his back using a surface receiving coil for the body. The following programs are used for the bile ducts investigation. A) Localizer (sighting image); B) T2–COR 2D FIESTA FATSAT (breath holding) - for more precise setting of slices for the MRCG program; C) 3D MRCG program without holding one’s breath, but using the 3D MRCG Rtr ASSET navigator-the program is synchronized with the movement of the diaphragm. The main parameters of the MRCG programs are shown in Table 2.

**Table 1** - Main parameters of the MDCT scanning programs

Parameters	Arterial phase	Portal phase	Delayed phase (phase of the hepatic veins)
Level	Liver hilum	Liver hilum	Liver hilum
Scan delay: Duration of study phases, automatic bolus tracking	15-20 sec 120 +HU	30-40sec	50-60sec
Tube voltage (kV)	120	120	120
Effective mas/ Quality ref mas	140	140	140
Rotation time	0.5s	0.5s	0.5s

**Table 2.** The main parameters of the MRCG programs

Parameters	COR 2D FIESTA FATSAT	3D MRCG Rtr ASSET
Layer	1	1
Quantity of slices	15-20	40-50
Diet factor	0	50%
FOV	300-330 mm	380 mm
FOV phase	100%	100%
Slice thickness	3-4.5 mm	1.5 mm
TR	1240 m/s	1800 m/s
TE	87 m/s	681 m/s
Averaging	1	1
Flip angel	150 deg.	170 deg.
Fat sat (with fat suppressed)	+	-

## RESULTS AND DISCUSSION

Up to and during conducting this research, 16 (18.4%) of the 88 potential examined donors have been operated on to collect a liver fragment. In RSSPMCS named after acad.V. Vakhidov, 4 liver transplants were performed, and the rest were operated on at Yashoda Hospital (India). Depending on the anthropometric data of the recipient, the volume of the transplant and the resected share of the donor were determined. Resection of the right lobe of the liver in order to obtain a transplant from the right lobe was performed in 12 (75%) donors and in 4 (25%) the left lobe of the liver was used as a transplant. The vascular and biliary anatomy of the liver and the mass of the resected liver fragment were also examined during the surgery.

During the surgery performed in donors it was revealed that in 8 (66.6%) from 12 who were resected the right lobe of the liver, the source of arterial circulation was own hepatic artery (OHA) extending from the celiac trunk, which corresponded to type I according to MDCT angiography and was identified in 7 (58.3%) donors at the preoperative stage. In 3 donors (25%) we revealed a discharge of the right hepatic artery from the common hepatic one which corresponds to type II at MDCT and was detected in 4 (33.3%) donors (8.3%) we registered a discharge of the right hepatic artery (RHA) from the superior mesenteric artery (SMA), which corresponded to type IV at MDCT.

In 4 donors the left lobe of the liver was used as a transplant. Intraoperatively, it was revealed that in 3 (75%) donors the left lobe was supplied from the OHA basin, which corresponded to type I according to MDCT and was detected in 3 (75%) donors. There was an abnormality of the left hepatic artery (LPA) from the total hepatic artery (OPA) in 1 donor (25%) which corresponds to type II according to MDCT and completely coincided with preoperative data. In all operated 16 donors blood supply of segment IV of the liver was also evaluated intraoperatively. Moreover, in 10 (62.5 ± 5.2%) of them blood supply was carried out due to the left lobar hepatic artery. Blood supply of the segment IV from the right hepatic artery was detected in 4 (25±4.1%) donors and in 2 (12.5±3.5%) a mixed type was revealed (left and right hepatic arteries). All donors were performed 3D reconstruction of the arterial bed of the liver (Figure 1). When evaluating the portal vein in 14 (87.5±3.6%) of the 16 operated donors, type I of portal vein (PV) branching (classical) was revealed. Trifurcation of the portal vein was detected in 1 donor (6.25 ± 2.4%) and in 1 donor at MDCT it was described as trifurcation, but type I was intraoperatively detected due to the close separation of the branch of the liver segment VI from bifurcation. 3D models of the liver portal bed were also reconstructed (Figure 2).



**Figure 1.** 3D-reconstruction of the arterial bed of the liver and abdominal aorta



**Figure 2.** 3D-model of the of the liver portal bed

An intraoperative assessment of venous outflow from the liver revealed that type I was detected in 9 (56.3%) patients - this is a separate inflow of the right hepatic vein (RHV), the median hepatic vein (MHV) and the left hepatic vein (LHV) into the postcava. In 4 cases (25%), type II of PV branching was observed, it was when MHV and LHV fell into a single mouth and type III, fusion of MHV and LHV into a single trunk was observed in 3 (18.7%) donors. In 12 donors who performed right-sided liver resection, RHV fell into postcava as a single trunk. Along with RHV in 2 donors (12.5%), an additional right vein from the segment VIII of the liver independently flowing into postcava was revealed. In 1 donor (6.3%), an additional lower right hepatic vein from the segment V to MHV was noted, which was reflected in the MDCT protocol. In 7 donors (43.7%), small additional lower right hepatic veins flowing into postcava were revealed intraoperatively. These small branches were hemodynamically insignificant and were ligated during the resection. 3D models of venous outflow of the liver were also used for volumetry of the liver (Figure 3).




**Figure 3.** 3D-reconstruction of the liver venous outflow


**The transplant volume**

When weighing during the operation the obtained transplant, the mass of the right lobe ranged from 560-1420 grams, the mass of the left lobe – from 240 to 670 grams. When calculating the volume of the transplant, absolute coincidence with intraoperative data was observed in 3 donors (18.7%) of the right lobe and in 1 donor (6.2%) of the left lobe of the liver. For the purpose of additional analysis of the obtained data, axial images obtained during MDCT angiography in the Digital Imaging and Communications in Medicine (DICOM) format were processed using the Vitrea (version 7.4.0.462, Vital Images) software, which allowed us to build virtual objects of the zones of interest to us and to study, based on 3D maps, the relative position of the vascular structures of the liver, as well as visually assess the future fragment of the liver and calculate the volume of the future transplant. Examples of reconstructed 3D maps are shown in [Figures 4 and 5](#).

Comparative data of MRCG with intraoperative ones were as follows. It was revealed that diagnosed type I on MRCG was detected in 13 (82%) from 16 operated patients. In 1 donor (6.2%), type I was determined on MRCG before the surgery, but type II was detected during surgery, which was explained by the close parallel lining of two right segmental ducts. In 1 donor (6.2%), 3 mouths of the bile ducts were revealed which corresponded to type III and was explained by the close fusion of two branches of the right anterior bile ducts to confluence. And in 1 (6.2%) donor we revealed 2 mouths of the transplant of the liver left lobe. Diagnostic informativity and accuracy of MDCT angiography in determining options and types of arterial and portal blood supply to the liver, as well as venous outflow pathways are presented in [Table 3](#) (P<0.05). Comparison of MDCT angiography and intraoperative data showed that the average error for the right lobe was 100±21.8 grams, for the left lobe 42.2±5.3 grams (P<0.05; [Table 4](#)). Diagnostic informativity and accuracy of MR cholangiography in determining biliary ducts are presented in [Table 5](#) (P<0.05).



**Figure 4.** 3D-model of the liver in the frontal projection



**Figure 5.** A 3D-model of the left lobe of the liver

**Table 3.** Diagnostic informativity of MDCT angiography in comparison with intraoperative data

Rates of the method informativity	Arterial blood supply of the liver	Variant of portal blood supply to the	Main ways of venous outflow
Sensitivity %	96.8	97.1	95.9
Accuracy %	92.5	93.7	91.3

**Citation:** Nazirov FG, Djuraeva NM, Vakhidova NT, Omonov OA and Salimov UR. Diagnostic informativity of the volume MDCT-angiography and MR-cholangiography in the pre- and intraoperative periods for the examination of donors of a liver fragment. *J. Life Sci. Biomed.*, 2019; 9(6): 151-156; [www.jlsb.science-line.com](http://www.jlsb.science-line.com)



**Table 4.** The average mass of the liver fragment (transplant) obtained on MDCT angiography and during surgery

Liver fragments	The average mass of the alleged transplant according to MDCT, grams	The average mass of the liver fragment obtained intraoperatively, grams
Right lobe	811.7±30.5	788.8±27.2
Left lobe	327.5±31.3	369±37.8

**Table 5.** Diagnostic informativity of MR cholangiography vs. intraoperative data

Rates of method informativity	Right lobar duct	Left lobar duct
Sensitivity %	96.8	97.1
Accuracy %	92.5	93.3

## CONCLUSION

According to preoperative MDCT angiography, the most common variant anatomy of the arterial blood supply to the liver was type Michels [31] classification, type I portal blood circulation according to Nakamura et al. [31] and type I venous outflow from the liver according to Soyer [32], which were determined intraoperatively at 95.7 % cases ( $P < 0.05$ ). MDCT angiography and MRCG in the definition of variant anatomy were 97.8% and 96.8%, respectively. The volume (mass) of the planned liver transplant obtained with MDCT volumetry was confirmed with intraoperative data in 92.8% ( $P < 0.05$ ). The data obtained indicate the high information content and the importance of these methods in planning liver transplantation.

## DECLARATIONS

### Acknowledgements

This work was supported by Republican Specialized Scientific and Practical Medical Center of Surgery named after academician V.Vakhidov, Tashkent, Uzbekistan

### Authors' Contributions

All authors contributed equally to this work.

### Competing interests

The authors declare that they have no competing interests.

## REFERENCES

1. Gauthier C.B. Konstantinov B.A. Tsirolnikova O.M. 2008. Liver Transplantation: A Guide for Physicians: 15-91.
2. Gauthier C.B., Tsirolnikova O.M., Filin A.B. et al. Experience of 25 transplantations of the right lobe of the liver from a living related donor. *Annals of the Russian Scientific Center of Medical Sciences*. 2002 (11): 30-36.
3. Kim E.F. Clinical and surgical aspects of intravital donation of liver fragments. Abstract. Dis. Dr. med. Science. 2008: 15-131.
4. Caruso S, Miraglia R, Maruzzelli L, Gruttadauria S, Luca A, Gridelli B. Imaging in liver transplantation. *World J Gastroenterol* 2009 (15): 675-83. [Google Scholar](#)
5. Kitaev V.M., Belova I.B., Kitaev C.B. Contrast agent for CT. 2008. *Computed tomography for liver diseases*. 1 (1.3): 17.
6. Adel El-Badrawy, Mahmoud El-Nahas, Asia El-Hendaw. Preoperative evaluation of living on or for liver transplantation using 64 MDCT // © *Journal of Medicine and Biomedical Sciences*, August, 2010, ISSN: P. 2073-2078.
7. Gautier S.V., Tsirolnikova O.M., Moysyuk Y.G., Akhaladze D.G., Tsirolnikova I.E., Silina O.V., Khizroev K.M., Monakhov A.R., Chekletsova E.V., Pets V.A., Poptsov V.N. Liver transplantation in children: six-year experience analysis. *Russian Journal of Transplantology and Artificial Organs*. 2014;16 (3): 54-62. [Google Scholar](#)
8. Kodama Y, Ng CS, Wu TT, Ayers GD, Curley SA, Abdalla EK, et al. 2007. Comparison of CT methods for determining the fat content of the liver. *AJR Am J Roentgenol*. 188:1307. [Google Scholar](#)
9. Ma X, Holalkere NS, Kambadakone RA, Mino-Kenudson M, Hahn PF, Sahani DV. 2009. Imaging-based quantification of hepatic fat: Methods and clinical applications. *Radiographics*, 29 (1253): 77. [Google Scholar](#)
10. Strasberg SM, Belghiti J, Clavien PA, Gadzijev E, Garden JO, Lau WY, et al. 2000. The Brisbane 2000 terminology of liver anatomy and resection. Terminology committee of the International Hepato-Pancreatico-Biliary Association. *HPB*. 2 (333):9. [Google Scholar](#)

11. Balakhnin P.V., Tarazov P.G., Polikarpov P.A. et al. 2004. Variants of arterial anatomy of the liver according to 1511 angiographies. *Ann. chir. Hepatol.* 9 (2): 14-21. [Google Scholar](#)
12. Nazirov F.G., Akbarov M.M., Saydazimov E.M., Nishanov M.Sh., Khakimov Yu.U. 2014. The use of modern technology for anatomical resections of the liver. *Bulletin of the Association of Physicians of Uzbekistan.* 4: 28-31.
13. Couinaud C. *Liver anatomy: portal or biliary segmentation.* Paris, 1957: 320. [Google Scholar](#)
14. Nazirov F.G., Akbarov M.M., Nishanov M.Sh., Saydazimov E.M., Khakimov Yu.U. 2013. Ways to improve the results of extensive anatomical liver resections. *Congress material "Actual problems of surgical hepatology". Donetsk:* 37-38.
15. Egorov V.I., Yashina N.I., Karmazanovsky G.G., Fedorov A.B. CT-angiography as a reliable method for verifying diseases, structural options, and postoperative changes in the arterial celiac-mesenteric basin. *Medical imaging 2009 No. 3* -p. 82.
16. Semenov A.V. *Related donors for liver transplantation: selection, examination, surgical tactics. /Abstract. of PhD dis.-M., 2003 - p. 21.*
17. Catalano OA, Singh AH, Uppot RN, Hahn PF, Ferrone CR, Sahani DV. Vascular and biliary variants in the liver: Implications for liver surgery. *Radiographics* 2008; 28 (359): 78. [Google Scholar](#)
18. Li K.W., Wen T.F., Yan L.N., Li B., Zeng Y., Zhao J.C., Wang W.T., Xu M.Q., Yang J.Y., Ma Y.K., Chen Z.Y., Huang B. Donor right hepatectomy in living donor liver transplantation: report of 143 cases. *Division of Liver Transplantation, West China Hospital of Sichuan University, Chengdu 610041, Sichuan Province, China. Hepatogastroenterology.* 2010 Sep-Oct; 57(102-103): 1232-6. [Google Scholar](#)
19. Tanaka K, Yamada T. Living donor liver transplantation in Japan and Kyoto University: What can we learn? *J Hepatol* 2005; 42: 25-8. [Google Scholar](#)
20. Zhuang ZG, Qian LJ, Gong HX, Zhou Y, Chai WM, Li QG, et al. Multidetector computed tomography angiography in the evaluation of potential living donors for liver transplantation: Single-center experience in China. *Transplant Proc* 2008; 40:2466-77. [Prime PubMed \(Link\)](#)
21. Sandrikov V.A., Semenov A.V., Khovrin V.V. Kim S.Yu., Burmistrov D.S., Korotneva N.A. Bogdanov- Berezovsky A.A. Possibilities of MR-cholangiography in the preoperative assessment of the variant structure of the bile ducts of liver fragments in living donors. *Annals of Surgical Hepatology.* (4): 57.
22. Kim KW, Lee J, Lee H, Jeong WK, Won HJ, Shin YM, et al. 2010. Right lobe estimated blood-free weight for liver transplantation: Accuracy of automated blood-free CT volumetry--preliminary results. *Radiology.* 256:433-40. [Google Scholar](#)
23. Radtke A, Sotiropoulos G.C., Nadalin S., Molmenti E.P., Schroeder T., Saner F.H., Sgourakis G., Cicinnati V.R., Valentin-Gamazo C., Broelsch C.E., Malago M. Preoperative volume prediction in adult live donor liver transplanttransplantation: 3-D volumetry approach to prevent miscalculations *Eur. J. Med. Res.* 2008 Jul 28; 13 (7): 319-26. [Google Scholar](#)
24. Wang F, Pan KT, Chu SY, Chan KM, Chou HS, Wu TJ, et al. Preoperative estimation of the liver graft weight in adult right lobe living donor liver transplantation using maximal portal vein diameters. *Liver Transpl* 2011; 17:373-80. Lindenbraten L.D. *Computed tomography Medical Radiology* 2006, 92-98. [Google Scholar](#)
25. Lee SW, Park SH, Kim KW, Choi EK, Shin YM, Kim PN, et al. Unenhanced CT for assessment of macrovesicular hepatic steatosis in living liver donors: Comparison of visual grading with liver attenuation index. *Radiology* 2007;244:479-85. [Google Scholar](#)
26. Krotova O.A., Granov D.A., Polysalov V.N., Pirtskhalava T.L., Borovik V.V., Rutkin I.O., Generalov M.I., Maistrenko D.N. Planning of surgical interventions on the liver based on the results of multilayer spiral computed tomography. *Annals of Surgical Hepatology.* 2010;15 (2): 31-37.
27. Khovrin V.V., Kim S.Yu., Kim E.F., Semenov A.V. Evaluation of variant vascular anatomy in potential living related donors of a liver fragment according to magnetic resonance and multispiral computed tomography. *Medical imaging.* 2010. 5. 37-43.
28. Lim JS, Kim MJ, Myoung S, Park MS, Choi JY, Choi JS, Kim SI. MR cholangiography for evaluation of hilar branching anatomy in transplantation of the right hepatic lobe from a living donor. *American Journal of Roentgenology.* 2008 Aug;191(2):537-45.. [Google Scholar](#)
29. Limanond P, Raman SS, Lassman C, Sayre J, Ghobrial RM, Busuttill RW, et al. Macrovesicular hepatic steatosis in living related liver donors: Correlation between CT and histologic findings. *Radiology* 2004; 230:276-80. [Google Scholar](#)
30. Michels NA. Newer anatomy of the liver and its variant blood supply and collateral circulation. *Am J Surg.* 1966; 112(3):337-347. doi: 10.1016/0002-9610(66)90201-7. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
31. Nakamura T, Tanaka K, Kiuchi T, Kashara M, Oike F, Ueda M, et al. Anatomical variants and surgical strategies in right lobe liver donor liver transplantation: Lessons from 120 cases. *Transplantation* 2002; 73:1896-903. [Google Scholar](#)
32. Soyer P, Levesque M, Legmann P, Fajadet P. Magnetic resonance imaging of malignant hepatic tumors in adults. *J Radiol.* 1992; 73(4):219-27. (PMID:1322459) [Article in French].